

# Policy on Early Childhood Caries (ECC): Consequences and Preventive Strategies

## Latest Revision

2025

**How to Cite:** American Academy of Pediatric Dentistry. Policy on early childhood caries (ECC): Consequences and preventive strategies. The Reference Manual of Pediatric Dentistry. Chicago, IL: American Academy of Pediatric Dentistry; 2025:96-100.

## Purpose

Early childhood caries (ECC), formerly referred to as nursing bottle caries and baby bottle tooth decay, remains a significant chronic disease of childhood and public health problem.<sup>1</sup> The American Academy of Pediatric Dentistry (AAPD) encourages health care professionals and caregivers to implement preventive practices that can decrease a child's risks of developing this preventable disease to reduce the burden on the child, the family, and society.

## Methods

This policy, initially developed by the American Academy of Pedodontics (now known as the AAPD) in collaboration with the American Academy of Pediatrics and adopted in 1978,<sup>2</sup> was last revised by the AAPD in 2021.<sup>3</sup> This update used an electronic search of English written articles in the dental and medical literature within the last 10 years, using the search terms (*evidence based dentistry* [Majr] OR *pediatric dentistry* [Majr] OR *dental care for children* [Majr]) AND (*dental caries* [Majr] OR *dental caries susceptibility* [Majr] OR *early childhood caries* [Tiab] OR *infant oral health* [Tiab] OR *oral microbiome* [Tiab] OR *ECC prevention* [Tiab] OR *patient well-being* [Tiab] OR *nutrition* [Tiab] OR *quality of life* [Majr] OR *ECC* [Tiab]). Two hundred forty-eight articles were identified in the search. Thirty-six articles were selected for review from this search and from references within the selected papers. When information from these articles did not appear sufficient or was inconclusive, expert and consensus opinions by experienced researchers and clinicians were considered.

## Background

In 1978, the American Academy of Pedodontics and the American Academy of Pediatrics released a joint statement, *Nursing Bottle Caries*, to address a severe form of caries associated with bottle usage.<sup>2</sup> Initial policy recommendations were limited to feeding habits, concluding that nursing bottle caries could be avoided if bottle feedings were discontinued soon after the first birthday.<sup>2</sup> An early policy revision added ad libitum breastfeeding as a causative factor.<sup>4</sup> Over the next 2 decades, however, recognizing that ECC was not solely associated with poor feeding practices, the AAPD adopted the term ECC to better reflect its multifactorial etiology. Contributors to ECC include<sup>5-9</sup> developmental (eg, enamel hypoplasia and hypomineralization), biological (eg, cariogenic oral microbiome

—especially Mutans streptococci [MS] and fungi), and behavioral (eg, poor oral hygiene, high-frequency sugar consumption) factors. Additionally, social and structural considerations (eg, socioeconomic status) have an overarching influence on the development of ECC.<sup>10</sup>

ECC is defined as “the presence of one or more decayed (noncavitated or cavitated lesions), missing (due to caries), or filled tooth surfaces in any primary tooth” in a child less than 72 months of age.<sup>11</sup> The definition of severe early childhood caries (S-ECC) is 1) any sign of smooth-surface caries in a child younger than 3 years of age; 2) from ages 3 through 5 years, 1 or more cavitated, missing (secondary to caries), or filled smooth surfaces in primary maxillary anterior teeth; or 3) a decayed, missing, or filled score of greater than or equal to 4 (age 3), greater than or equal to 5 (age 4), or greater than or equal to 6 (age 5).<sup>11</sup>

Epidemiologic data from a 2013-2018 national health survey clearly indicate that ECC remains highly prevalent in poor and near-poor US preschool children.<sup>12</sup> For the overall population of preschool children, the prevalence of ECC, as measured by decayed and filled tooth surfaces (dfs), is unchanged from previous surveys.<sup>12</sup> However, the proportion of children with severe ECC (S-ECC) increased significantly.<sup>12</sup> The consequences of ECC include a higher risk of new caries lesions in both the primary and permanent dentitions,<sup>13,14</sup> hospitalizations and emergency room visits,<sup>15</sup> high treatment costs,<sup>16</sup> loss of school days,<sup>17,18</sup> diminished ability to learn,<sup>18</sup> nutritional deficiencies,<sup>19-21</sup> and reduced<sup>22-24</sup> oral health-related quality of life.

Traditional microbial risk markers for ECC include acidogenic-aciduric bacterial species, particularly MS.<sup>25</sup> Studies using direct culture with arbitrarily primed polymerase chain reaction (AP-PCR) fingerprinting and other traditional techniques have shown that MS may be transmitted vertically from parent or caregiver to child and horizontally from other individuals in the immediate environment.<sup>26,27</sup> Advanced technologies that include DNA and RNA sequencing (ie, next

## ABBREVIATIONS

**AAPD:** American Academy of Pediatric Dentistry. **CWF:** Community water fluoridation. **ECC:** Early childhood caries. **Majr:** MeSH major topic. **MeSH:** Medical subject heading. **MS:** Mutans streptococci. **Tiab:** Title and abstract.

generation sequencing), along with machine learning and artificial intelligence, reveal the complexity of the oral microbiome and have highlighted other bacterial species (eg, *Scardovia wiggsiae*, *Veillonella* spp.) and fungi (eg, *Candida* species) that also may be associated with ECC.<sup>28-31</sup> In addition to vertical and horizontal transmission, studies on the development of the oral microbiome since birth support the importance of healthy diets and environmental exposures.<sup>32,33</sup> It is recommended that parental education and counseling on the importance of a healthy microbiome and diet in infancy be conducted as early as possible as the oral microbiome associated with ECC can be influenced<sup>16,7,29,34,35</sup> by the child's sex, genetic variants in taste, socioeconomic factors, dietary practices, and lifestyle behaviors.

An associated risk factor to microbial etiology is high consumption of sugars.<sup>36</sup> Bottle feeding, particularly nighttime bottle feeding with sweet drinks or sugar-sweetened beverages<sup>7</sup> (eg, juice, fruit-flavored drinks), continuous use of a sippy or no-spill cup containing sugar-sweetened beverages, and frequent in-between meal consumption of sugar-sweetened snacks or drinks increase the risk of caries.<sup>37</sup> Although benefits of breastfeeding are clear,<sup>38</sup> frequent (on demand) or nocturnal breastfeeding beyond the first year of life is associated with ECC.<sup>39-42</sup> A 2019 consensus statement via the American Academy of Pediatrics, American Heart Association, Academy of Nutrition and Dietetics, and AAPD recommends that sugar in foods and drinks should be avoided in children under 2 years of age.<sup>43</sup> Additional recommendations include complete avoidance of juice in children younger than 1 year and no more than 4 ounces of 100% fruit juice a day for children between the ages of 1 and 3 years.<sup>43,44</sup>

Community water fluoridation (CWF) as a primary prevention method is considered a key strategy for preventing dental caries.<sup>45</sup> Children with lifetime exposure to CWF show significantly lower dental caries levels than those without, with the benefit being most pronounced in primary teeth.<sup>46</sup> In addition to reducing the prevalence of severe caries, the use of CWF in high-risk populations may reduce caries-related visits and help avoid preventable dental surgery under general anesthesia.<sup>47</sup> CWF has multiple benefits and attenuates income-related inequalities in dental caries in the US.<sup>48</sup> Despite conflicting reports<sup>49-51</sup> of an association between fluoride consumption and a child's intelligence quotient (IQ), apart from an increased incidence of enamel fluorosis, the literature fails to provide credible evidence linking CWF with negative health outcome in the US at the currently regulated CWF level (0.7 ppm).<sup>52-54</sup>

Current best practice to reduce the risk of ECC includes twice-daily brushing with 1000 ppm fluoridated toothpaste for all children in optimally-fluoridated and fluoride-deficient communities.<sup>55-57</sup> When determining the risk-benefit of fluoride, the key issue is mild fluorosis versus preventing dental disease. A smear or rice-sized amount of 1000 ppm fluoridated toothpaste (approximately 0.1 mg fluoride; see Figure) should be used for children younger than 3 years of




Figure. Comparison of a smear (left) with a pea-sized (right) amount of toothpaste.

age. A pea-sized amount of 1000 ppm fluoridated toothpaste (approximately 0.25 mg fluoride) is appropriate for children aged 3 to 6 years.<sup>58</sup> Parents can ensure their child uses an appropriate amount of toothpaste by dispensing it onto a soft, age-appropriate sized toothbrush and performing or assisting with toothbrushing of preschool-aged children. Minimal or no rinsing can maximize the beneficial effect of fluoride in the toothpaste.<sup>59</sup> Less than twice daily toothbrushing and difficulties in performing the procedure during the preschool years were significant determinants of caries prevalence at the age of 5 years.<sup>36</sup>

Professionally-applied topical fluoride treatments are efficacious in reducing the incidence of ECC. The recommended professionally-applied fluoride treatment for children at risk for ECC who are younger than 6 years is 5% sodium fluoride varnish (NaFV; 22,500 ppm fluoride).<sup>60-62</sup> Evidence suggests that preventive interventions within the first year of life are critical.<sup>63</sup> For this reason, establishment of a dental home within 6 months of the eruption of the first tooth and no later than 12 months of age is especially important in populations at risk for ECC. This may be implemented with the help of medical professionals who, in many cases, are being trained to provide oral screenings, apply preventive measures, counsel caregivers, and refer infants and toddlers for dental care.<sup>64</sup> Additionally, 38% silver diammine fluoride (SDF) is effective for the arrest of cavitated caries lesions in primary teeth<sup>65,66</sup> and may be used as part of an ongoing caries management plan within the context of a dental home.

### Policy statement

The AAPD recognizes ECC as a significant chronic disease resulting from an imbalance of multiple risk and protective factors over time. The AAPD encourages the following evidence-based preventive measures to decrease the risk of developing ECC and to promote optimal oral health for infants and children.

- Establishment of a dental home within 6 months of eruption of the first tooth and no later than 12 months of age to conduct caries-risk assessment and parental education.
- At the initial dental visit and periodic dental appointments, delivery of anticipatory guidance including home

oral hygiene instructions (eg, parental brushing with an age-appropriate sized toothbrush and volume of fluoridated toothpaste) and dietary counseling that emphasizes

- avoidance of on-demand or nocturnal baby bottle- and breastfeeding beyond 12 months of age.
  - encouragement for children between 6 and 12 months old to drink 4 to 8 ounces of water per day and to abstain from all juice consumption.
  - avoidance of foods and drink with added sugars in children under 2 years of age and restriction of 100% fresh or reconstituted juice to no more than 4 ounces a day for children between the ages of 1 and 3 years.
  - avoidance of frequent consumption of sugar-containing liquids and/or solid foods.
- Professional application of fluoride varnish for children at risk for ECC.

Additionally, the AAPD supports interprofessional education and public health efforts to promote oral prevention, including

- working with medical health care professionals to ensure all infants and toddlers have access to dental screenings, counseling, and preventive procedures with a consistent, unified message.
- supporting CWF as a primary prevention for dental caries to reach underserved and vulnerable communities.
- educating legislators, policy makers, and third-party payors regarding the consequences of and preventive strategies for ECC, emphasizing the importance of access to care for all.
- advocating for reimbursement systems to allow access for all children and educational reforms that emphasize evidence-based preventive and comprehensive management of ECC.

## References

1. American Academy of Pediatric Dentistry. Proceedings of the conference: Innovations in the Prevention and Management of Early Childhood Caries. October 2014. Chicago, IL. *Pediatr Dent* 2015;37(3):198-299.
2. American Academy of Pedodontics, American Academy of Pediatrics. Nursing bottle caries. January 1978. Reference Manual 1991-1992. Chicago, IL: American Academy of Pediatric Dentistry; 1991:27.
3. American Academy of Pediatric Dentistry. Policy on early childhood caries (ECC): Consequences and preventive strategies. The Reference Manual of Pediatric Dentistry. Chicago, IL: American Academy of Pediatric Dentistry; 2021;81-4.
4. American Academy of Pediatric Dentistry. Baby bottle tooth decay, revised 1993. *Pediatr Dent* 1993;15(7):27.
5. Americano GCA, Jacobsen PE, Soviero VM, Haubek D. A systematic review on the association between molar incisor hypomineralization and dental caries. *Int J Paediatr Dent* 2017;27(1):1-78.
6. de Jesus VC, Shikder R, Oryniak D, et al. Sex-based diverse plaque microbiota in children with severe caries. *J Dent Res* 2020;99(6):703-12.
7. Khan SY, Schroth RJ, Cruz de Jesus V, et al. A systematic review of caries risk in children <6 years of age. *Int J Paediatr Dent* 2024;34(4):410-31. Available at: “<https://onlinelibrary.wiley.com/doi/10.1111/ipd.13140>”. Accessed July 20, 2025.
8. Lu Y, Lin Y, Li M, He J. Roles of *Streptococcus mutans-Candida albicans* interaction in early childhood caries: A literature review. *Front Cell Infect Microbiol* 2023;13:1-7.
9. Tinanoff N. Introduction to the conference: Innovations in the Prevention and Management of Early Childhood Caries. *Pediatr Dent* 2015;37(4):198-9.
10. American Academy of Pediatric Dentistry. Policy on social determinants of oral health and oral health disparities. The Reference Manual of Pediatric Dentistry. Chicago, IL: American Academy of Pediatric Dentistry; 2025:47-51.
11. Drury TF, Horowitz AM, Ismail AI, et al. Diagnosing and reporting early childhood caries for research purposes. *J Public Health Dent* 1999;59(3):192-7.
12. Kotha A, Vemulapalli A, Mandapati SR, Aryal S. Prevalence, trends, and severity of early childhood caries in the United States: National health and nutritional examination survey data 2013 to 2018. *Pediatr Dent* 2022;44(4):261-8.
13. Lam PPY, Chua H, Ekamaram M, Lo ECM, You CKY. Does early childhood caries increase caries development among school children and adolescents? A systematic review and meta-analysis. *Int J Environ Res Public Health* 2022;19(20):7-12.
14. Schroth RJ, Rothney J, Sturym M, et al. A systematic review to inform the development of a Canadian caries risk assessment tool for use by primary healthcare providers. *Int J Paediatr Dent* 2021;31(6):767-91
15. Musa AAR, Sethi S, Poirier BF, Oliver KJ, Jensen ED. Non-traumatic emergency department dental visits among patients 0-25 years of age: A systematic review and meta-analysis. *Int J. Paediatr Dent* 2023;33(5):457-67.
16. Owens PL, Manski RJ, Weiss AJ. Emergency Department Visits Involving Dental Conditions, 2018. Healthcare Cost and Utilization Project, Statistical Brief #280. Agency for Healthcare Research and Quality; August 2021. Available at: “<https://hcup-us.ahrq.gov/reports/stat-briefs/sb280-Dental-ED-Visits-2018.pdf>”. Accessed August 7, 2025.
17. Edelstein BL, Reisine S. Fifty-one million: A mythical number that matters. *J Am Dent Assoc* 2015;146(8):565-6.
18. Ruff RR, Senthil S, Susser SR, Tsutsui A. Oral health, academic performance, and school absenteeism in children and adolescents. *J Am Dent Assoc* 2019;150(2):111-21.

19. Carvalho Silva C, Mendes R, Manso MDC, Gavinha S, Melo P. Prenatal or childhood serum levels of vitamin D and dental caries in paediatric patients: A systematic review. *Oral Health Prev Dent* 2020;18(4):653-67.
20. Easwaran HN, Annadurai A, Muthu MS, et al. Early childhood caries and iron deficiency anaemia: A systematic review and meta-analysis. *Caries Res* 2022;56(1):36-46.
21. Schroth RJ, Pierce A, Rodd C, et al. Improvement in serum vitamin D following dental rehabilitation to treat severe early childhood caries. *Pediatr Dent* 2022;44(4):278-83.
22. Farsi DJ, Farsi NJ, El-Housseiny AA, Turkistani JM, Farsi NM. Impact of dental rehabilitation on oral health-related quality-of-life in healthy children and those with special health care needs. *J Contemp Dent Pract* 2018;19(4):367-74.
23. Grant CG, Daymont C, Rodd C, et al. Oral health-related quality of life of Canadian preschoolers with severe caries after dental rehabilitation under general anesthesia. *Pediatr Dent* 2019;41(3):221-8.
24. Raghu R, Gauba K, Goyal A, Kapur A, Supta A, Singh SK. Oral health-related quality of life of children with early childhood caries before and after receiving complete oral rehabilitation under general anesthesia. *Int J Clin Pediatr Dent* 2021;14(Suppl 2):S117-S123.
25. Kopycka-Kedzierawski DT, Billings RJ, Feng C, et al. A prospective longitudinal study of early childhood caries onset in initially caries-free children. *JDR Clin Trans Res* 2023;8(4):394-401.
26. Doméjean S, Zhan L, DenBesten PK, Stamper J, Boyce WT, Featherstone JD. Horizontal transmission of mutans streptococci in children. *J Dent Res* 2010;89(1):51-5.
27. Manchanda S, Sardana D, Liu P, et al. Horizontal transmission of streptococcus mutans in children and its association with dental caries: A systematic review and meta-analysis. *Pediatr Dent* 2021;43(1):E1-E12.
28. Hahishengallis E, Parsaei Y, Klein MI, Koo H. Advances in the microbial etiology and pathogenesis of early childhood caries. *Mol Oral Microbiol* 2017;32(1):24-34.
29. Khan MW, de Jesus VC, Mittermuller BA, et al. Role of socioeconomic factors and interkingdom crosstalk in the dental plaque microbiome in early childhood caries. *Cell Rep* 2024;43(8):1-17. Available at: "[https://www.cell.com/cell-reports/pdf/S2211-1247\(24\)00985-9.pdf](https://www.cell.com/cell-reports/pdf/S2211-1247(24)00985-9.pdf)". Accessed August 7, 2025.
30. Li Y, Tanner A. Effect of antimicrobial interactions on the oral microbiota associated with early childhood caries. *Pediatr Dent* 2015;37(3):226-44.
31. Mira A. Oral microbiome studies: Potential diagnostic and therapeutic implications. *Adv Dent Res* 2018;29(1):71-7.
32. Dashper SG, Mitchel HL, Lê Cao KA, et al. Temporal development of the oral microbiome and prediction of early childhood caries. *Sci Rep* 2019;9(1):19732. Available at: "<https://doi.org/10.1038/s41598-019-56233-0>". Accessed August 7, 2025.
33. Dzidic M, Collado MC, Abrahamsson T, et al. Oral microbiome development during childhood: An ecological succession influenced by postnatal factors and associated with tooth decay. *ISME J* 2018;12(9):2292-306.
34. de Jesus VC, Mittermuller BA, Hu P, Schroth RJ, Chelikani P. Genetic variants in taste genes play a role in oral microbial composition and severe early childhood caries. *iScience* 2022;25(12):105489.
35. Kopycka-Kedzierawski DT, Scott-Anne K, Ragusa PG, et al. Social, psychological, and behavioral predictors of salivary bacteria, yeast in caries-free children. *JDR Clin Trans Res* 2022;7(2):163-73.
36. Moynihan PJ, Kelly SAM. Effect on caries of restricting sugars intake: Systematic review to inform WHO guidelines. *J Dent Res* 2014;93(1):8-18.
37. Tinanoff NT, Kanellis MJ, Vargas CM. Current understanding of the epidemiology, mechanism, and prevention of dental caries in preschool children. *Pediatr Dent* 2002;24(6):543-51.
38. Salone LR, Vann WF, Dee DL. Breastfeeding: An overview of oral and general health benefits. *J Am Dent Assoc* 2013;144(2):143-51.
39. Panchanadikar NT, Abirami S, Muthu MS, Selvakumar H, Jayakumar P, Agarwal A. Breastfeeding and its association with early childhood caries – An umbrella review. *J Clin Pediatr Dent* 2022;46(2):75-85.
40. Peres KG, Chaffee BW, Feldens CA. Breastfeeding and oral health: Evidence and methodological challenges. *J Dent Res* 2018;97(3):251-8.
41. Shrestha SK, Arora A, Manohar N, Ekanayake K, Foster J. Association of breastfeeding and early childhood caries: A systematic review and meta-analysis. *Nutrients* 2024;16(9):1355.
42. Wong PD, Birken CS, Parkin PC, et al. Total breast-feeding duration and dental caries in healthy urban children. *Acad Pediatr* 2017;17(3):310-5.
43. Lott M, Callahan E, Welker Duffy E, Story M, Daniels S. Healthy Beverage Consumption in Early Childhood: Recommendations from Key National Health and Nutrition Organizations. Consensus Statement. Durham, NC: Healthy Eating Research; 2019. Available at: "<https://healthyeatingresearch.org/wp-content/uploads/2019/09/HER-HealthyBeverage-ConsensusStatement.pdf>". Accessed August 7, 2025.
44. Heyman MB, Abrams SA; American Academy of Pediatrics Committee on Nutrition. Fruit juice in infants, children, and adolescents: Current recommendations. *Pediatrics* 2017;139(6):e20170967.
45. Iheozor-Ejiofor Z, Walsh T, Lewis SR, et al. Water fluoridation for the prevention of dental caries. *Cochrane Database Syst Rev* 2024;10(10):CD010856. Available at: "<https://doi.org/10.1002/14651858.CD010856.pub2>". Accessed August 7, 2025.

*References continued on the next page.*

46. Slade GD, Grider WB, Maas WR, Sanders AE. Water fluoridation and dental caries in U.S. children and adolescents. *J Dent Res* 2018;97(10):1122-8.
47. Lee HH, Faundez MA, LoSasso AT. A cross-sectional analysis of community water fluoridation and prevalence of pediatric dental surgery among Medicaid enrollees. *JAMA Network Open* 2020;3(8):e205882.
48. Sanders AE, Grider WB, Mass WR, Curiel JA, Slade GD. Association between water fluoridation and income-related dental caries of U.S. children and adolescents. *JAMA Pediatr* 2019;173(3):288-90.
49. Broadbent JM, Thomson WM, Ramrakha S, et al. Community water fluoridation and intelligence: Prospective study in New Zealand. *Am J Public Health* 2015;105(1):72-6.
50. Green R, Lanphear B, Hornung R, et al. Association between maternal fluoride exposure during pregnancy and IQ scores in offspring in Canada. *JAMA Pediatr* 2019;173(10):940-8. Available at: "<https://jamanetwork.com/journals/jamapediatrics/fullarticle/2748634>". Accessed August 7, 2025.
51. Taylor KW, Eftim SE, Sibrizzi CA, et al. Fluoride exposure and children's IQ scores: A systematic review and meta-analysis. *JAMA Pediatr* 2025;179(3):282-92.
52. Centers for Disease Control and Prevention. CDC Scientific Statement on Community Water Fluoridation. May 2024. Available at: "<https://www.cdc.gov/fluoridation/about/statement-on-the-evidence-supporting-the-safety-and-effectiveness-of-community-water-fluoridation.html>". Accessed August 7, 2025.
53. Levy SM. Caution needed in interpreting the evidence base on fluoride and IQ. *JAMA Pediatr* 2025;179(3):231-4.
54. Tinanoff N, Dhar V, Caffrey E. Community water fluoridation: Evidence of efficacy and risks. *Pediatr Dent* 2024;46(6):370-1.
55. American Dental Association Council on Scientific Affairs. Fluoride toothpaste use for young children. *J Am Dent Assoc* 2014;145(2):190-1.
56. Boustedt K, Dahlgren J, Twetman S, Roswall J. Toothbrushing habits and prevalence of early childhood caries: A prospective cohort study. *Eur Arch Paediatr Dent* 2020;21(1):155-9.
57. Santos AP, Oliveira BH, Nadanovsky P. Effects of low and standard fluoride toothpastes on caries and fluorosis: Systematic review and meta-analysis. *Caries Res* 2013;47(5):382-90.
58. Wright JT, Hanson N, Ristic H, Whall CW, Estrich CG, Zentz RR. Fluoride toothpaste efficacy and safety in children younger than 6 years. *J Am Dent Assoc* 2014;145(2):182-9.
59. Sjögren K, Birkhed D. Factors related to fluoride retention after toothbrushing and possible connection to caries activity. *Caries Res* 1993;27(6):474-7.
60. American Academy of Pediatric Dentistry. Fluoride therapy. *The Reference Manual of Pediatric Dentistry*. Chicago, IL: American Academy of Pediatric Dentistry; 2025:372-8.
61. American Academy of Pediatric Dentistry. Policy on use of fluoride. *The Reference Manual of Pediatric Dentistry*. Chicago, IL: American Academy of Pediatric Dentistry; 2025:110-2.
62. Weyant RJ, Tracy SL, Anselmo T, Beltrán-Aguilar EJ, Donly KJ, Frese WA. Topical fluoride for caries prevention: Executive summary of the updated clinical recommendations and supporting systematic review. *J Am Dent Assoc* 2013;144(11):1279-91.
63. Lee JY, Bouwens TJ, Savage MF, Vann WF. Examining the cost-effectiveness of early dental visits. *Pediatr Dent* 2006;28(2):102-5, discussion 192-8.
64. Douglass AB, Douglass JM, Krol DM. Educating pediatricians and family physicians in children's oral health. *Academic Pediatr* 2009;9(6):452-6.
65. Crystal YO, Marghalani AA, Ureles SD, et al. Use of silver diamine fluoride for dental caries management in children and adolescents, including those with special health care needs. *Pediatr Dent* 2017;39(5):E135-E145.
66. Gao SS, Zhao IS, Hiraishi N, et al. Clinical trials of SDF in arresting caries among children: A systematic review. *JDR Clin Trans Res* 2016;1(3):201-10.